



Eur päisch s Patentamt
European Patent Office
Office européen des brevets



Publication number: **0 341 947 B1**

(12)

EUROPEAN PATENT SPECIFICATION

- (45) Date of publication of patent specification: **31.08.94** (51) Int. Cl.⁵: **C11D 3/39, C11D 3/386, C12N 15/00**
- (21) Application number: **89304620.1**
- (22) Date of filing: **08.05.89**

(54) Enzymatic detergent and bleaching composition.

(30) Priority: **09.05.88 GB 8810954**

(43) Date of publication of application:
15.11.89 Bulletin 89/46

(45) Publication of the grant of the patent:
31.08.94 Bulletin 94/35

(84) Designated Contracting States:
CH DE ES FR GB IT LI NL SE

(56) References cited:
EP-A- 0 271 152
LU-A- 65 030

(73) Proprietor: **UNILEVER PLC**
Unilever House
Blackfriars
London EC4P 4BQ (GB)

(84) Designated Contracting States:
GB

(73) Proprietor: **UNILEVER N.V.7**
Weena 455
NL-3013 AL Rotterdam (NL)

(84) Designated Contracting States:
CH DE ES FR IT LI NL SE

(72) Inventor: **Lagerwaard, Cornelis Abraham**
Liguster 5
NL-3224 TM Hellevoetsluis (NL)
Inventor: **Cornelissen, Johannes Mattheus**
Platanendreef 27
NL-3137 CN Vlaardingen (NL)
Inventor: **Swarthoff, Ton**
Debussystraat 3
NL-3223 TE Hellevoetsluis (NL)
Inventor: **Klugkist, Jan**
Staverenhoeve 12
NL-3137 GN Vlaardingen (NL)
Inventor: **Thom, David**
Cherry Bank Manorial Road
Parkgate Wirral Merseyside (GB)

(74) Representative: **Kan, Jacob Hendrik, Dr. et al**
Unilever N.V.
Patent Division
P.O. Box 137
NL-3130 AC Vlaardingen (NL)

Not : Within nine months from the publication of the mention of the grant of the European patent, any person may give notice to the European Patent Office of opposition to the European patent granted. Notice of opposition shall be filed in a written reasoned statement. It shall not be deemed to have been filed until the opposition fee has been paid (Art. 99(1) European patent convention).

Description

The present invention relates to an enzymatic detergent and bleaching composition comprising as essential ingredients a lipolytic enzyme and a bleaching system.

Detergent compositions comprising lipase are known. For example GB 1 372 034 (Unilever) discloses lipase from *Pseudomonas* in specific nonionic-containing detergent compositions for soaking fabrics.

USP 3 950 277 (Procter & Gamble) also describes fabric-soaking compositions: the described compositions comprise lipase and lipase activators and a number of lipases from microorganism and other sources are mentioned: those particularly mentioned as preferred are Amano CE, Amano M-AP, Takeda 1969-4-9, and Meito MY-30 lipases, but no indications are given of the form in which the lipase is to be prepared or used.

USP 4 011 169/NL 74 08763 (Procter & Gamble) describes the use of a similar range of enzymes in the preparation of additives for washing agents (detergent compositions).

Examples of known lipase-containing detergent compositions are provided by EP 0 205 208 and 0 206 390 (Unilever), which relate to lipases related to those from *Ps. fluorescens*, *P. gladioli* and *Chromobacter* in detergent compositions.

EP 0 214 761 (Novo) and EP 0 258 068 (Novo), each give detailed description of lipases from certain microorganisms, and also give certain uses in detergent additives and detergent compositions for the enzymes described. EP 0 214 761 gives detailed description of lipases derived from organisms of the species *Pseudomonas cepacia*, and certain uses therefor. EP 0 258 068 gives detailed description of lipases derived from organisms of the genus *Thermomyces/Humicola*, and certain uses therefor.

Also believed to be in use in certain areas is a lipase-containing granular detergent composition containing about 37% detergent actives including 5% nonionic detergent and the remainder substantially anionic detergent, about 16% zeolite, about 60 LU/g lipase, plus protease and other normal detergent additives.

Further examples of known lipase-containing detergent compositions are provided by JA 63-078000 (1988) (Lion Corp/K Mukoyama et al) which discloses properties and uses of a *Pseudomonas* lipase, including use in a lipase-containing system based on 10-40 % surfactant (e.g., sodium C14-C18 alpha-olefin sulphonate), as well as other conventional detergent ingredients.

In EP 0 268 456 (Clorox), there is described in connexion with Table 10(b) an experimental washing solution containing lipase and about 1 microgram/ml sodium dodecyl sulphate.

In US Patent No 4,707,291 (hereby incorporated by reference herein), detergent compositions have been described which contain a special class of lipolytic enzymes. These compositions may also contain a bleaching agent such as sodium perborate, either as such or in admixture with a low temperature bleach activator, e.g., tetraacetyl-ethylene-diamine (TAED). The lipases mentioned in US Patent No 4,707,291 are significantly less affected by a bleaching system than other lipases. These bleaching systems comprise sodium perborate and TAED.

We have now surprisingly found that a certain type of lipase, defined herein below, is quite compatible with bleaching systems which are stronger than the sodium perborate/TAED system, such systems being defined in more detail hereafter. This type of lipase is more stable in the presence of these bleaching systems than some of the lipases mentioned US Patent No 4,707,291 and is in particular more stable than the lipases mentioned in US Patent No 4,707,291 in the presence of proteases in these bleaching systems.

The lipase used in the present invention is a lipase produced by cloning, by rDNA technique, the gene encoding the lipase produced by the fungus *Humicola lanuginosa* and expressing the gene in *Aspergillus oryzae* as host. This product is manufactured and commercially available from Novo Industri A/S, Denmark, under the trade name Lipolase (see 'Biotechnology Newswatch', published 7th March 1988, page 6). In the relevant manufacturer's technical leaflet on Lipolase (Trade Mark) it is said that this lipase is not stable in the presence of e.g., hypochlorite as bleaching agent.

Equivalence of a lipase enzyme produced by rDNA technique to the lipase defined above can be determined in practice for example by an immunological reaction of identity between the two lipase materials and/or by determination of substantial aminoacid sequence homology therebetween. In some such equivalent lipases, of which the use is within the scope of the invention, the pattern of glycosylation of the lipase protein differs.

EP 0 305 216 (Novo) also describes suitable lipase enzymes for use in the practice of this invention, and is hereby incorporated by reference herein.

The lipase used in the present invention is included in the detergent and bleaching composition in such an amount that the final composition has a lipolytic enzyme activity of from 100 to 0.005 LU/mg, preferably 25 to 0.05 LU/mg of the composition.

A Lipase Unit (LU) is that amount of lipase which produces 1/micromol of titratable fatty acid per minute in a pH stat. under the following conditions: temperature 30 °C; pH = 9.0; substrate is an emulsion of 3.3 wt.% of olive oil and 3.3% gum arabic, in the presence of 13 mmol/l Ca^{++} ; and 20 mmol/l NaCl in 5 mmol/l Tris-Buffer. An alternative definition of the lipase unit is given in EP 0 258 068 (Novo), which mentions (inter alia) lipases suitable for use in the practice of this invention and is hereby incorporated by reference.

Naturally, mixtures of the above lipase with other known lipases can be used. The lipase(s) can be used in its (their) non-purified form or in a purified form, e.g., purified with the aid of well-known adsorption methods, such as phenyl sepharose adsorption techniques.

The bleaching system used according to the present invention is stronger than the sodium perborate/TAED system. This latter system, through a perhydrolysis reaction, forms a peroxyacid, i.e., peracetic acid, but at a rather low rate. The bleaching systems according to the present invention must be stronger than this sodium perborate/TAED system, by which is to be understood that the system either is based on a peracid (inorganic or organic) which is stronger than the peracetic acid or yields, on perhydrolysis, an organic peracid, including peracetic acid, faster than the sodium perborate/TAED system. The bleaching system may consist of a bleaching agent as such or may consist of a bleaching agent together with a bleach precursor. As bleaching agent as such alkali metal monopersulphates, furthermore organic peracids such as diperoxy dodecanedioic acid, diperoxy tetradecanedioic acid, diperoxyhexadecane dioic acid, mono- and diperazelaic acid, mono- and diperbrassylic acid, monoperoxy phthalic acid, perbenzoic acid, can be used, either as acid or in the form of their salts.

When a system comprising a bleach precursor is used, this system comprises a bleaching agent which reacts with a bleach precursor to form a peracid in solution faster than the sodium perborate/TAED system. By faster is meant that the precursor will have a rate of peroxy acid release of at least 2 (two) times, preferably at least 5 (five) times faster than TAED under the same conditions.

Typical examples of such systems are sodium perborate with sodium nonanoyloxy benzene sulphonate or sodium trimethyl hexanoyloxy benzene sulphonate or sodium acetoxy benzene sulphonate or sodium benzoyloxy benzene sulphonate.

The preferred systems of the present invention are sodium perborate with sodium nonanoyloxy benzene sulphonate, diperoxy dodecane dioic acid or monopersulphate.

In general, the amount of the bleaching system in the composition varies from 1-50%, usually from 5-40% by weight. When a bleach precursor is present, the molar ratio of the bleach precursor to the percompound such as sodium perborate varies from 1:1 to 1:35, preferably from 1:2 to 1:20. Mixtures of various bleaching agents and various bleach precursors in accordance with the invention can also be used.

The compositions of the present invention may furthermore contain one or more detergent active materials, such as soaps, anionic, nonionic, cationic and zwitterionic synthetic detergents or mixtures thereof. Usually the amount of detergent active material present in the composition will range from 1-50%, preferably 2-40% and particularly preferably 5-30% by weight. Suitable examples of detergent active materials can be found in Schwartz, Perry and Berch "Surface Active Agents and Detergents", Vol I (1949) and Vol II (1958) and M Schick "Nonionic Surfactants" Vol I (1967).

The compositions may furthermore include the usual detergent ingredients in the usual amounts. They may be unbuilt or built, and may be of the zero-P type (i.e., not containing phosphorus-containing builders). Thus, the compositions may contain from 1-60%, preferably from 5-30% by weight of one or more organic and/or inorganic builders. Typical examples of such builders are the alkali metal ortho-, pyro- and tri-polyphosphates, alkali metal carbonates, either alone or in admixture with calcite, alkali metal citrates, alkali metal nitrilotriacetates, carboxymethyloxy succinates, zeolites, polyacetal carboxylates and so on.

The compositions may furthermore comprise lather boosters, foam depressors, anti-corrosion agents, soil-suspending agents, sequestering agents, anti-soil redeposition agents, perfumes, dyes, stabilizing agents for the enzymes and bleaching agents and so on. They may also comprise enzymes other than lipases, such as proteases, amylases, oxidases and cellulases. In this respect it has been found that, whereas proteases, are often affected by strong bleaches, in the present invention, when used together with the lipases of the present invention, the overall performance of the enzyme system is often not significantly affected.

In general, the compositions may comprise such other enzymes in an amount of 0.01-10% by weight. For proteases, the amount, expressed in proteolytic activity, is usually from 0.1-50 GU/mg based on the final composition.

A GU is a glycine unit, which is the amount of proteolytic enzyme which under standard incubation conditions produces an amount of terminal NH_2 -groups equivalent to 1 microgram/ml of glycine.

The compositions of the present invention can be formulated in any desired form, such as powders, bars, pastes, liquids, etc.

The invention will further be illustrated by way of Example. The invention described herein extends to modifications and variations as will be apparent to the skilled reader, and combinations and subcombinations of the features mentioned herein.

EXAMPLE 1

The stability of various lipases in the presence of a bleaching system was measured as follows:

To a solution of 4 g/l of a detergent composition* and 0.03 g/l Dequest 2041 in water with a hardness of 30 ° FH and a temperature of 30 ° C, an amount of lipase is added to obtain 15-20 lipase units/ ml.

The pH is adjusted with NaOH to pH 10.0 at 30°C. At t=0 a bleach system is added:

- a) 292 mg/l TAED (65% pure) and 700 mg/l sodium perborate monohydrate or
- b) 1880 mg/l DPDA (12% pure) or
- c) 822 mg/l SNOBS (80% pure) and 1500 mg/l sodium perborate monohydrate or
- d) 506 gm/l MPS (in the form of the commercial product Caroate R) or
- e) 475 mg/l P15 (95% pure) and 700 mg/l sodium perborate monohydrate.

This yields 1.5 mmolar peracid in solution for all bleach systems. The lipase stability is measured by determining the residual lipase activity with the pH-stat. method.

- Dequest 2041 = ethylene diamine tetra (methylene phosphonic acid)
- TAED = tetraacetyl ethylene diamine
- DPDA = diperoxy dodecanedioic acid
- SNOBS = sodium nonanoyloxy benzene sulphonate
- MPS = sodium monopersulphate
- P15 = sodium benzoyloxy benzene sulphonate

* The detergent composition had the following formulation:

	% by weight
Sodium dodecyl benzene sulphonate	6.5
C14-C15 primary alcohol, condensed with 11 moles of ethylene oxide	2.0
Sodium stearate	1.0
Sodium silicate	7.0
Sodium carboxymethyl cellulose	0.5
Na ₂ SO ₄	37.0
Pentasodium triphosphate	15.0
Trisodium orthophosphate	5.0
Fluorescer	0.2
Ethylene diamine tetraacetic acid	0.5
Water	6.2
Dyes	0.01

The following results were obtained :

Lipase ex.	Trade-name	No bleach activity ¹⁾		TAED/perb. activity ¹⁾		SNORS/perb. activity ¹⁾		DPDA activity ¹⁾		MPS activity ¹⁾		PLS activity ¹⁾	
		10	30 t1/2	10	30 t1/2	10	30 t1/2	10	30 t1/2	10	30 t1/2	10	30 t1/2
		min	min(min)	min	min(min)	min	min(min)	min	min(min)	min	min(min)	min	min(min)
Humicola lanuginosa	Lipolase	100	99 >30	100	85 >30	95	83 >30	95	70 >30	94	72 >30	100	96 >30
Pseudomonas gladioli	-	97	97 >30	97	92 >30	90	80 >30	100	99 >30	89	80 >30	92	77 >30
Chromobacter viscosum	Toyo Jozo	100	95 >30	100	86 >30	80	58 >30	107	98 >30	85	62 >30	93	63 >30
Pseudomonas fluorescens	Amano P	100	98 >30	92	100 >30	85	80 >30	109	100 >30	89	85 >30	102	83 >30
Pseudomonas cepacia	ex NOVO	93	90 >30	98	94 >30	89	81 >30	100	98 >30	97	96 >30	95	84 >30
Aspergillus niger	Amano AP 6	110	90 >30	100	64 43	14	<5 2	62	<5 12	78	40 24	30	9 7
Mucor Niehei	SP 225	67	46 28	95	58 35	81	46 28	86	68 >30	13	5 4	73	44 26

1) residual lipase activity (% of input)
t1/2 = half life time

EXAMPLE 2

Example 1 was repeated, but now in the presence of 20 GU (glycine unit)/ml Savinase[®], a proteolytic enzyme ex NOVO.

The following results were obtained :

Lipase	No bleach activity ¹⁾		TAED/perb. activity ¹⁾		SNOBS/perb. activity ¹⁾		DPDA activity ¹⁾		MPS activity ¹⁾		PLS activity ¹⁾	
	10	30	10	30	10	30	10	30	10	30	10	30
	min	min	min	min	min	min	min	min	min	min	min	min
Lipolase	100	100	100	88	94	85	96	90	94	72	100	96
Ps. gladioli	81	56	76	51	76	55	90	70	63	47	81	42
Amano P	51	17	56	20	40	16	60	24	43	27	55	15
Ps. cepacia	65	35	72	38	65	34	59	42	54	32	65	28
Toyo Jozo	79	48	71	38	72	47	82	52	38	22	74	29
Amano AP6	96	83	82	38	45	3	61	15	91	79	55	24
Esterase MM	64	21	38	15	43	12	68	25	10	5	74	25

* in an experiment with Kazusase, a proteolytic enzyme ex Shova Denko, the following results were obtained: 95% 84% >30.

** With Alcalase instead of Savinase, the results were as follows: 97% 93% >30.

*** With Alcalase instead of Savinase, the results were as follows: 90% 76% >30.

**** With Kazusase instead of Savinase, the results were as follows: 99% 89% >30.

1) residual lipase activity (% of input)
t1/2 = half life time

Claims

1. A detergent composition comprising from 1-50% by weight of one or more detergent-active materials, from 0-60% by weight of a builder, from 1-50% by weight of a bleaching agent and lipolytic enzyme in

an amount of 0.005-100 lipolytic units per milligram of the composition, characterised in that the bleaching agent is based on an inorganic or organic peracid or salt thereof which is stronger than peracetic acid or comprises a bleaching agent and a bleach precursor which yields, on perhydrolysis, a peracid faster than the system comprising sodium perborate + tetraacetyl ethylene diamine, and the lipolytic enzyme is a lipase produced by cloning by rDNA technique the gene encoding the lipase produced by Humicola lanuginosa (syn. Thermomyces lanuginosus) and expressing the gene in Aspergillus oryzae as host.

2. A composition according to Claim 1, characterised in that the bleaching agent is an alkali metal persulphate.
3. A composition according to Claim 1 characterised in that the bleaching agent is selected from the group consisting of diperoxy dodecanedioic acid, diperoxy tetradecanedioic acid, diperoxyhexadecanedioic acid, mono- and diperazelaic acid, mono- and diperbrassylic acid, monoperoxy phthalic acid, perbenzoic acid, and their salts.
4. A composition according to Claim 1, characterized in that the bleaching agent comprises a bleaching agent and a bleach precursor which forms a peracid in solution at least two times faster than tetraacetyl ethylene diamine under the same conditions.
5. A composition according to Claim 4, characterised in that the bleaching agent comprises sodium perborate and a bleach precursor selected from the group consisting of sodium nonanoyloxy benzene sulphonate, sodium trimethyl hexanoyloxy benzene sulphonate, sodium acetoxy benzene sulphonate and sodium benzoyloxy benzene sulphonate.
6. A composition according to any of Claims 1-5, characterized in that it further contains a proteolytic enzyme in an amount of 0.1-50 GU/mg of the composition.
7. A composition according to any of Claims 1-6, characterised in that the lipase enzyme is Lipolase (Trade Mark).

Patentansprüche

1. Reinigungsmittelzusammensetzung, umfassend 1 bis 50 Gew.% eines oder mehrerer reinigungsmittelaktiver Materialien, 0 bis 60 Gew.% eines Aufbaustoffes, 1 bis 50 Gew.% eines Bleichmittels und ein lipolytisches Enzym in einer Menge von 0,005 bis 100 lipolytischen Einheiten pro Milligramm der Zusammensetzung, dadurch gekennzeichnet, daß das Bleichmittel auf einer anorganischen oder organischen Persäure oder einem Salz davon basiert, das stärker als Peressigsäure ist, oder ein Bleichmittel und eine Bleichmittelvorstufe umfaßt, die bei einer Perhydrolyse schneller zu einer Persäure führt als das Natriumperborat und Tetraacetyl ethylendiamin umfassende System, und daß das lipolytische Enzym aus einer Lipase besteht, die durch Klonen mit Hilfe einer rDNA-Technik des für die durch Humicola lanuginosa (Synonym Thermomyces lanuginosus) gebildete Lipase kodierenden Gens und Exprimieren des Gens in Aspergillus oryzae als Wirt hergestellt ist.
2. Zusammensetzung nach Anspruch 1, dadurch gekennzeichnet, daß das Bleichmittel aus einem Alkali-metallpersulfat besteht.
3. Zusammensetzung nach Anspruch 1, dadurch gekennzeichnet, daß das Bleichmittel aus der Gruppe Diperoxydodecandicarbonsäure, Diperoxytetradecandicarbonsäure, Diperoxyhexadecandicarbonsäure, Mono- und Diperazelaensäure, Mono- und Diperbrassyssäure, Monoperoxyphthalsäure, Perbenzoesäure und ihren Salzen ausgewählt ist.
4. Zusammensetzung nach Anspruch 1, dadurch gekennzeichnet, daß das Bleichmittel ein Bleichmittel und eine Bleichmittelvorstufe, die mindestens zweimal schneller als Tetraacetyl ethylendiamin unter denselben Bedingungen eine Persäure in Lösung bildet, umfaßt.
5. Zusammensetzung nach Anspruch 4, dadurch gekennzeichnet, daß das Bleichmittel Natriumperborat und eine Bleichmittelvorstufe, ausgewählt aus der Gruppe Natriumnonaoxyloxybenzolsulfonat, Natriumtri-

methylhexanoyloxybenzolsulfonat, Natriumacetoxybenzolsulfonat und Natriumbenzoyloxybenzolsulfonat, umfaßt.

6. Zusammensetzung nach einem der Ansprüche 1 bis 5, dadurch gekennzeichnet, daß sie des weiteren ein proteolytisches Enzym in einer Menge von 0,1 bis 50 GU/mg der Zusammensetzung enthält.
7. Zusammensetzung nach einem der Ansprüche 1 bis 6, dadurch gekennzeichnet, daß das Lipaseenzym aus Lipolase (Handelsbezeichnung) besteht.

10 Revendications

1. Composition détergente comprenant de 1 à 50% en poids d'un ou plusieurs détergent(s) actifs, de 0 à 60% en poids d'un adjuvant, de 1 à 50% en poids d'un agent de blanchiment et une enzyme lipolytique en une quantité de 0,005 à 100 Unités lipolytiques par mg de la composition, caractérisée en ce que l'agent de blanchiment est à base d'un peracide minéral ou organique ou d'un sel de celui-ci plus fort que l'acide peracétique ou comprend un agent de blanchiment et un précurseur de blanchiment donnant, lors d'une perhydrolyse, un peracide plus rapidement que le système comprenant le perborate de sodium plus la tétraacétyléthylènediamine et l'enzyme lipolytique est une lipase produite par clonage par la technique de l'ADNr du gène encodant la lipase produite par Humicola lanuginosa (syn. Thermomyces lanuginosus) et expression du gène dans Aspergillus oryzae à titre d'hôte.
2. Composition selon la revendication 1, caractérisée en ce que l'agent de blanchiment est un persulfate de métal alcalin.
3. Composition selon la revendication 1, caractérisée en ce que l'agent de blanchiment est choisi parmi l'acide diperoxy-dodecanedioïque, l'acide diperoxytetradecanedioïque, l'acide diperoxyhexadecanedioïque, les acides mono- et diperazélaïques, les acides mono- et diperbrassyliques, l'acide monoperoxyphthalique, l'acide perbenzoïque et leurs sels.
4. Composition selon la revendication 1, caractérisée en ce que l'agent de blanchiment comprend un agent de blanchiment et un précurseur de blanchiment formant un peracide en solution au moins deux fois plus vite que la tétra-acétyléthylènediamine dans les mêmes conditions.
5. Composition selon la revendication 4, caractérisée en ce que l'agent de blanchiment comprend du perborate de sodium et un précurseur de blanchiment choisi parmi le nonanoyloxybenzènesulfonate de sodium, le triméthylhexanoyloxybenzènesulfonate de sodium, l'acétoxy-benzènesulfonate de sodium et le benzoyloxybenzènesulfonate de sodium.
6. Composition selon la revendication 1 à 5, caractérisée en ce qu'elle contient en outre une enzyme protéolytique en une quantité de 0,1 à 50 UG/mg de composition.
7. Composition selon l'une quelconque des revendications 1 à 6, caractérisée en ce que l'enzyme lipase est la Lipolase (marque déposée).